

FUSE Observations of the Post-AGB Star ZNG 1 in the Globular Cluster M5 (NGC 5904)

W. Van Dyke Dixon

*Department of Physics & Astronomy, The Johns Hopkins University,
 3400 N. Charles Street, Baltimore, MD 21218, USA*

Thomas M. Brown

*Space Telescope Science Institute, 3700 San Martin Drive, Baltimore,
 MD 21218, USA*

Wayne B. Landsman

*Science Systems and Applications, Inc., Code 681, NASA Goddard
 Space Flight Center, Greenbelt, MD 20771, USA*

Abstract. We have observed the hot post-AGB star ZNG 1 in the globular cluster M5 with the *Far Ultraviolet Spectroscopic Explorer (FUSE)*. From the resulting spectrum, we derive an effective temperature $T_{\text{eff}} \sim 45,000$ K, a rotational velocity $v_{\text{rot}} \sim 100$ km s $^{-1}$, carbon and nitrogen abundances approximately ten times solar, a wind velocity $v_{\infty} \sim 1000$ km s $^{-1}$, and evidence for an expanding shell of material around the star. The carbon and nitrogen enhancements suggest dredge-up of nuclear-processed material on the AGB. The high rotational velocity may reflect a previous merger with a binary companion.

As part of a project to study mixing and mass loss in the atmospheres of asymptotic-giant-branch (AGB) stars, we have observed the post-AGB star ZNG 1 (Zinn, Newell, & Gibson 1972) in M5 (NGC 5904) with the *Far Ultraviolet Spectroscopic Explorer (FUSE)*. The resulting spectrum, presented in Fig. 1, suggests that the star has experienced both processes – and that the mass loss continues.

The upper two panels show narrow absorption features of H I and C II blueshifted by ~ 190 km s $^{-1}$ relative to the star. C III absorption at the same velocity is also present, but is not shown here. These features may represent an intermediate-velocity cloud or a shell of material previously ejected by the star. Broad P-Cygni profiles can be seen in both components of the O VI doublet (middle panel), each punctuated by a discrete absorption component (DAC) blueshifted by 920 km s $^{-1}$ relative to the star. The redder O VI doublet is interstellar. Photospheric features of nitrogen and carbon can be seen in the bottom panel. Their redshift (60 km s $^{-1}$ relative to the ISM) is consistent with that of the cluster (Harris 1996). The star's iron abundance is close to the cluster mean ($[\text{Fe}/\text{H}] = -1.29$; Harris 1986), but its carbon and nitrogen abundances are ten times solar, suggesting that products of helium burning were mixed to the surface while the star was on the AGB. The star's photospheric features are




Figure 1. Far-UV spectrum of ZNG 1 in M5 obtained with *FUSE*. The top and middle panels present evidence for a high-velocity stellar wind and expanding shell. In the bottom panel, a synthetic spectrum (generated with the non-LTE code of Hubeny & Lanz 1995) with $T_{\text{eff}} = 45,000$ K, $\log g = 4.5$, $[\text{Fe}/\text{H}] = -1.3$, and carbon and nitrogen abundances ten times solar is plotted over the data. A higher nitrogen abundance is inconsistent with other features not shown here.

well fit by a rotational velocity of $\sim 100 \text{ km s}^{-1}$, remarkably fast for a post-AGB star. The star may have been spun up in a merger with a binary companion.

References

- Harris, W. E. 1996, *AJ*, 112, 1487
 Hubeny, I. & Lanz, T. 1995, *ApJ*, 439, 875
 Zinn, R. J., Newell, E. B., & Gibson, J. B. 1972, *A&A*, 18, 390